

# The influence of age and metformin treatment status on reported gastrointestinal side effects with liraglutide treatment in type 2 diabetes

Thong, K.y.; Gupta, P. Sen; Blann, Andrew; Ryder, R.e.j.

DOI:

[10.1016/j.diabres.2015.04.009](https://doi.org/10.1016/j.diabres.2015.04.009)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Thong, KY, Gupta, PS, Blann, A & Ryder, REJ 2015, 'The influence of age and metformin treatment status on reported gastrointestinal side effects with liraglutide treatment in type 2 diabetes', *Diabetes Research and Clinical Practice*, vol. 109, no. 1, pp. 124-129. <https://doi.org/10.1016/j.diabres.2015.04.009>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication in *Diabetes Research and Clinical Practice*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Diabetes Research and Clinical Practice*, Vol 109, Issue 1, July 2015. DOI: 10.1016/j.diabres.2015.04.009

Eligibility for repository checked

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Accepted Manuscript

Title: The influence of age and metformin treatment status on reported gastrointestinal side effects with liraglutide treatment in type 2 diabetes

Author: K.Y. Thong P.S. Gupta A.D. Blann R.E.J. Ryder



PII: S0168-8227(15)00191-6  
DOI: <http://dx.doi.org/doi:10.1016/j.diabres.2015.04.009>  
Reference: DIAB 6354

To appear in: *Diabetes Research and Clinical Practice*

Received date: 21-1-2015  
Revised date: 23-3-2015  
Accepted date: 14-4-2015

Please cite this article as: K.Y. Thong, P.S. Gupta, A.D. Blann, R.E.J. Ryder, The influence of age and metformin treatment status on reported gastrointestinal side effects with liraglutide treatment in type 2 diabetes, *Diabetes Research and Clinical Practice* (2015), <http://dx.doi.org/10.1016/j.diabres.2015.04.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- Gastrointestinal side effects with liraglutide increases with patient age
- Patients not on metformin had more significant gastrointestinal side effects
- Renal function was not an independent predictor of gastrointestinal side effects

Original Research Article – Clinical Research

**The influence of age and metformin treatment status on reported gastrointestinal side effects with liraglutide treatment in type 2 diabetes**

K Y Thong<sup>a</sup>, P S Gupta<sup>b</sup>, A D Blann<sup>c</sup>, R E J Ryder<sup>b</sup>

<sup>a</sup>School of Medicine and Pharmacology, University of Western Australia, Perth, Australia

<sup>b</sup>Diabetes Centre, City Hospital, Birmingham, United Kingdom

<sup>c</sup>University of Birmingham Centre for Cardiovascular Science, Department of Medicine, City Hospital, Birmingham, United Kingdom

Corresponding author:

Dr Ken Yan Thong  
Department of Diabetes and Endocrinology,  
Rockingham General Hospital,  
Elanora Drive,  
Rockingham, WA 6168  
Australia

Telephone: +618 95994697  
Fax: +618 95994737  
Email: [kythong@gmail.com](mailto:kythong@gmail.com)

Keywords: GLP-1, liraglutide, metformin, gastrointestinal side effects

## ABSTRACT

### Aim

Treatment of type 2 diabetes with glucagon-like peptide-1 (GLP-1) receptor agonists may be limited by gastrointestinal side effects (GISE) in some patients. Risk factors for developing GISE are not known. We analysed patient characteristics that were associated with GISE among patients treated with the GLP-1 receptor agonist liraglutide.

### Methods

Data was obtained from an audit database of liraglutide use based in clinical practice in the UK. Patients were grouped into those who did not report GISE, those who reported GISE but continued liraglutide and those who discontinued liraglutide due to GISE within 26 weeks of treatment. Baseline variables of age, diabetes duration, HbA<sub>1c</sub>, weight, BMI, blood pressure, lipids, gender, ethnicity, alanine transaminotransferase, estimated glomerular filtration rate (eGFR) and diabetes treatment types were tested for possible associations with GISE outcome. Significant variables in univariate analyses were entered into ordinal logistic regression analyses.

### Results

4442 patients were suitable for analysis. 3905 (87.9%) did not report GISE, 297 (6.7%) and 240 (5.4%) had GISE and continued and discontinued treatment, respectively. Age, weight, eGFR, metformin status and insulin status were associated with GISE outcome in univariate analyses ( $P$  all  $<0.05$ ). In the final regression model, age (adjusted OR 1.15 [95%CI 1.05,1.26],  $P = 0.002$ ) and non-metformin use (adjusted OR 0.76 [95%CI 0.60,0.96],  $P = 0.020$ ) were associated with worse GISE outcome.

### Conclusion

Older age and non-metformin use were associated with more significant GISE leading to discontinuation of liraglutide treatment. The reasons for these findings are unclear and warrant further investigation.

## 1. Introduction

Glucagon-like peptide 1 (GLP-1) secretion by intestinal L cells leads to an increase in insulin secretion, suppression of glucagon secretion, slowing of gastric emptying and suppression of appetite. However, endogenous GLP-1 is rapidly degraded by the dipeptidyl peptidase-4 (DPP-4) enzyme. The use of GLP-1 receptor agonists which are resistant to such degradation or drugs inhibiting of the DPP-4 enzyme, forms the pharmacological basis of incretin-based therapies in patients with type 2 diabetes [1].

Liraglutide, a once-daily injected GLP-1 receptor agonist helps improve glycaemia and reduce weight in patients with type 2 diabetes [2]. However, a disadvantage of the GLP-1 receptor agonist drug class including that of liraglutide is the occurrence of gastrointestinal side effects (GISE). Common symptoms include nausea, vomiting or diarrhoea [1]. Liraglutide has been shown to have less persistent GISE than exenatide twice daily, the first available GLP-1 receptor agonist [3]. The side effects of GLP-1 receptor agonists also tend to improve with longer duration of treatment, possibly due to the development of tachyphylaxis [4]. However, there remains to be patients who discontinue drug treatment due to severe or persistent symptoms; between four to five percent of patients treated with liraglutide 1.2 mg once daily discontinued liraglutide treatment due to GISE in the Liraglutide Effect and Action in Diabetes (LEAD) trials [5-7]. In the United Kingdom (UK), liraglutide 1.2 mg once daily is recommended for use by the National Institute for Health and Care Excellence (NICE) as a second or third line diabetes treatment after metformin and/or sulphonylurea in patients with body mass indices (BMI) greater than 35 kg/m<sup>2</sup> [8].

To date, there is a lack of large-scale data on the rates of GISE among patients treated with GLP-1 receptor agonists in real-life clinical practice. There is also little data on identifying patients who may be at risk. Using information from a nationwide audit database of liraglutide use in clinical practice, we investigated whether there were identifiable risk factors for developing GISE among patients treated with liraglutide.

## 2. Methods

### 2.1 Subjects

Data was obtained from the Association of British Clinical Diabetologists (ABCD) nationwide liraglutide audit database. ABCD had invited diabetes centres in hospitals and primary care across UK to submit anonymised data of patients treated with liraglutide in routine clinical practice during the period of 2009 to 2013. 117 diabetes centres enrolled and submitted varying degrees of data depending on date of participation, the frequency of patients' health visits and duration of liraglutide treatment that had taken place. Data entry and submission was performed using audit software provided by ABCD.

Data requested included patients' age, gender, ethnicity, pre- and post-liraglutide treatment information including diabetes treatments, glycated haemoglobin (HbA<sub>1c</sub>), body weight, BMI, blood pressure, lipid parameters, alanine transaminotransferase (ALT) and creatinine, whenever these data were available. Contributors were also asked to report on occurrence of any GISE or other possible treatment related adverse events, as well as the main reason for liraglutide discontinuation if this occurred.

6238 patients had baseline data sent to ABCD. For this study, we excluded patients without at least one follow-up data submission (n=1296) and those with less than 13 weeks data contribution or follow-up unless these patients were noted to have discontinued liraglutide (n=500). Remaining 4422 patients were used for analyses.

## 2.2 Outcome and risk variables

Patients were classified into three GISE outcome groups: a) patients without GISE, b) patients reporting GISE but continued with liraglutide treatment and c) patients who discontinued liraglutide due to GISE before 26 weeks (the three groups were coded 0, 1, and 2). We defined symptoms of GISE as those of nausea, diarrhoea, vomiting, crampy abdominal pain, constipation, belching, reflux, flatulence or similar related terms.

## 2.3 Statistical analyses

Univariate associations of all variables in the database were tested against GISE outcome using ANOVA, Kruskal-Wallis test and tabulated statistics depending on whether the variables were continuous and normally or non-normally distributed, or were categorical variables, respectively. Results shown for estimated glomerular filtration rates (eGFR) were those calculated by the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation and data on eGFR was first analysed as a continuous variable. Sensitivity analyses were performed using eGFR derived by the Cockcroft-Gault equation adjusted for body surface area using the DuBois formula, and the Modification of Diet in Renal Disease (MDRD) equation. All eGFR equations used are shown in Appendix 1 [9,10]. eGFR was also analysed as a categorical variable based on chronic kidney disease (CKD) groups: CKD group 1 (eGFR >90 ml/min/1.73m<sup>2</sup>), CKD Group 2 (eGFR 60-89 ml/min/1.73m<sup>2</sup>) and CKD Group 3 (eGFR 30-59 ml/min/1.73m<sup>2</sup>). The associations between baseline diabetes treatments and GISE were also tested. In particular, the association between metformin use and GISE was tested due to clinical trials of liraglutide generally showing higher rates of GISE among patients on metformin compared with patients on sulphonylurea [5,7,11]. The association between insulin use and GISE was tested due to our previous finding of more frequent reports of GISE among insulin-treated patients as compared with non-insulin-treated patients when started on exenatide twice daily [12]. Variables with statistically significant association in univariate analyses ( $P < 0.050$ ) were entered into an ordinal logistic regression analysis, with liraglutide dose (1.2 mg versus 1.8 mg, and prior exenatide use treated as covariates. Variables in logistic regression with the highest  $P$  value were sequentially removed until a final model with all variables with  $P < 0.10$  was achieved. Due to findings of age being a significant risk factor for GISE in the final regression model, the results of rates of GISE according to decades of age are also shown. Statistical analyses were performed using Minitab<sup>®</sup> Release 16 (Minitab Ltd, Coventry, UK).

## 3. Results

Baseline characteristics of the 4422 patient were, mean ( $\pm$ SD), age  $56 \pm 11$  years, HbA<sub>1c</sub>  $9.4 \pm 1.7\%$  ( $79 \pm 19$  mmol/mol), weight  $110.5 \pm 22.5$  kg and BMI  $38.8 \pm 7.2$  kg/m<sup>2</sup>. Median (inter-quartile range) diabetes duration was 9 [6-13] years. 41.2% of patients were on insulin at liraglutide initiation as opposed to the licensed prescribing indications for liraglutide.

537 (12.1%) patients reported GISE of which 297 (6.7%) continued liraglutide treatment while 240 (5.4%) discontinued liraglutide treatment due to GISE. The total number of patients discontinuing liraglutide within 26 weeks of treatment, including those who discontinued due to GISE, was 606 (13.6%).

Results of univariate analyses are shown in Table 1 and 2. Variables significantly associated with GISE were older age, lower weight, poorer renal function, the use of insulin and non-use of metformin ( $P$  all  $<0.05$ ).

The results of ordinal logistic regression analyses are shown in Table 3. In the final model of 4213 patients, a rise in one standard deviation of age which approximated an increase of 10 years of age (adjusted OR 1.15 [95%CI 1.05,1.27],  $P = 0.002$ ) and non-metformin use (adjusted OR for metformin versus non metformin use 0.76 [95%CI 0.60,0.96],  $P = 0.020$ ) were associated with GISE outcome. The proportions of patients reporting GISE according to decades of life are shown in Figure 1.

Sensitivity analyses performed on eGFR derived by the Cockcroft-Gault and the MDRD equations showed similar findings to that when eGFR was derived by the CKD-EPI equation; the association of eGFR with GISE outcome in univariate analyses remained significant, and the association in multivariate analyses remained non-significant. Analysing eGFR according to CKD groups also did not alter our univariate or multivariate findings.

## 4. Discussion

Our analysis of the data from the ABCD liraglutide audit database hopes to bring renewed focus on the issue of drug tolerability of GLP-1 receptor agonists, rather than drug efficacy *per se*. To our knowledge, our analysis may be the first published attempt to characterise risk factors for developing GISE among patients treated with this class of treatment. There is now an expanding array of diabetes treatment options for type 2 diabetes beyond metformin, including sulphonylureas, thiazolidinediones, insulin, GLP-1 receptor agonists, DPP-4 inhibitors and more recently sodium glucose transporters 2 inhibitors [13]. Knowledge on potential risk factors for GISE may help guide clinicians in whether to select GLP-1 receptor agonists as an option for treatment escalation. However, the results of our study should be mainly treated as hypothesis-generating due to the susceptibility to biases of data in an audit, as well as the post-hoc decision to analyse GISE data in such a manner.

Reported rates of nausea, vomiting or diarrhoea vary between GLP-1 receptor agonists but were generally lower for liraglutide or exenatide once weekly as compared with exenatide twice daily in head-to-head studies [3,13]. In the ABCD audit, the rates of all combined reported GISE with liraglutide 1.2 mg were low at 12.1%. Rates of liraglutide discontinuation due to GISE in the audit were 5.4% and were comparable with the rates of 4 to 5% in the LEAD trials [5-7].

Our results suggest that older age and non-metformin use are risk factors for GISE and drug discontinuation among patients treated with liraglutide. While eGFR was also significantly associated with GISE in univariate analysis, eGFR was highly collinear with age. Removing eGFR in the multivariate analysis resulted in the variable of age being significantly associated with GISE while eGFR itself did not achieve significance in any of our multivariate models. In increasing decades of life from age 30 years, there was a steady rise in reported rates of GISE (9.7% to 20.0%). Discordantly, there was also a high rate of reported GISE among patients age 20 to 29



years (15.6%) although the total number of patients were small in this group (64 patients). The reason for older age being a risk factor for more significant GISE is not clear. It is also possible that the higher rate of drug discontinuation in the elderly may signify a more precautionary approach to their management rather than them being more prone to developing severe GISE. Prescribing information for liraglutide suggests that no dose adjustment is required for patients older than 65 years, but highlighted the limited experience of the use of liraglutide among patients with age more than 75 years [14].

The findings of patients not on metformin reporting greater drug discontinuation due to GISE was initially a surprising finding. Clinical trials of liraglutide versus placebo or active comparator treatment have reported generally higher rates of GISE when there was background treatment with metformin [5,7] as compared with a trial of liraglutide added to glimepiride without background metformin use [10]. However, in the latter trial, glimepiride treatment was used as part of a forced treatment-titration rather than due to metformin intolerance. In clinical practice, metformin is well-recognised as the first line diabetes treatment; patients in the audit not on metformin were possibly those who had encountered side effects to metformin earlier in their treatment algorithm. However, the audit did not capture data that could verify this, this hence being a limitation of this study. There is a plausible hypothesis that patients may share intolerance to both metformin and GLP-1 receptor agonists. The side effect profiles for both treatments are similar. Metformin has been shown to increase GLP-1 levels after an oral glucose load in obese subjects with or without diabetes, and the rise in GLP-1 may be a possible cause of the anorectic effects of metformin [15].

We have previously reported more frequent discontinuation of liraglutide in the audit among patients with mild and moderate renal impairment, a similar finding in the current univariate analysis [16]. However, when adjusted for other variables including age, renal function became non-significantly associated with GISE outcomes. A pharmacokinetic study testing the 0.75 mg dose of liraglutide showed no increase in drug exposure in patients with renal impairment [17]. The Prescribing Information for liraglutide have reported that liraglutide is metabolised endogenously much like large proteins rather than having a specific organ as a major route of elimination, and with no significant elimination through the faeces or urine [14]. Hence our finding that poorer renal function was not an independent risk factor for GISE with liraglutide treatment would be consistent with the pharmacokinetic studies above.

The strengths of our study include the fact that all available variables were tested and data was obtained from a large number of centres across UK. Weaknesses of the study include the potential for loss of follow-up or lack of data input among patients who discontinued liraglutide early due to GISE, as well as the potential for the audit not capturing data on other unknown determinants of GISE.

In conclusion, the results from our analyses would suggest caution in using liraglutide in older patients due to poorer drug tolerability. It may also be worthwhile investigating whether patients susceptible to GISE caused by metformin treatment are also at greater risk of developing GISE to GLP-1 receptor agonists.

## Conflict of interest statement

KYT has received speaker fees and educational sponsorships from Novo Nordisk, manufacturer of liraglutide. PSG has received educational sponsorships from Novo Nordisk. REJR has received speaker fees, consultancy fees and educational sponsorships from Novo Nordisk. ADB has no conflicts of interest to declare.

### Acknowledgment

We thank all the nationwide contributors for submitting data on patients on liraglutide (listed in Appendix 2).

### REFERENCES

1. Neumiller JJ. Incretin-Based Therapies. *Med Clin North Am* 2015; 99: 107-29.
2. Blonde L, Russell-Jones D. The safety and efficacy of liraglutide with or without oral antidiabetic drug therapy in type 2 diabetes: an overview of the LEAD 1-5 studies. *Diabetes Obes Metab* 2009; 11: 26-34.
3. Buse JB, Rosenstock J, Sesti G, Schmidt WE, Montanya E, Brett JH, et al. Liraglutide once a day versus exenatide twice a day for type 2 diabetes: a 26-week randomised, parallel-group, multinational, open-label trial (LEAD-6). *Lancet* 2009; 374: 39-47.
4. Nauck MA, Kemmeries G, Holst JJ, Meier JJ. Rapid tachyphylaxis of the glucagon-like peptide 1-induced deceleration of gastric emptying in humans. *Diabetes*. 2011; 60: 1561-5.
5. Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, et al. Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes. *Diabetes Care* 2009; 32: 84-90.
6. Garber A, Henry R, Ratner R, . Liraglutide versus glimepiride monotherapy for type 2 diabetes (LEAD-3 Mono): a randomized, 52-week, phase III, double-blind, parallel-treatment trial. *Lancet* 2009; 373: 473-81.
7. Zinman B, Gerich J, Buse JB, Lewin A, Schwartz S, Raskin P, et al. Efficacy and safety of the human glucagon-like peptide-1 analog liraglutide in combination with metformin and thiazolidinedione in patients with type 2 diabetes (LEAD-4 MET+TZD). *Diabetes Care* 2009; 32: 1224-30.
8. NICE technology appraisal guidance 203. Liraglutide for the treatment of type 2 diabetes. 2010. Available at <http://guidance.nice.org.uk/TA203> (accessed 10 December 2014).
9. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3<sup>rd</sup>, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009; 150: 604-12.
10. Chudleigh RA, Dunseath G, Peter R, Harvey JN, Ollerton RL, Luzio S, et al. Influence of body weight on the performance of glomerular filtration rate estimators in subjects with type 2 diabetes. *Diabetes Care* 2008; 31: 47-9.
11. Marre M, Shaw J, Brändle M, Bebakar WMW, Kamaruddin NA, Strand J, et al. Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with type 2 diabetes (LEAD-1 SU). *Diabet Med* 2009; 26: 268-78.
12. Thong KY, Jose B, Sukumar N, Cull ML, Mills AP, Sathyapalan T, Shafiq, et al.. Safety, efficacy and tolerability of exenatide in combination with insulin in the Association of British Clinical Diabetologists nationwide exenatide audit. *Diabetes Obes Metab* 2011; 13: 703-10.
13. Inzucchi SE, Bergenstal RM, Buse JB, Diamant M, Ferrannini E, Nauck M, et al. Management of hyperglycaemia in type 2 diabetes: a patient-centred approach. Position

- statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2012; 55: 1577-96.
14. Drucker DJ, Buse JB, Taylor K, Kendall DM, Trautmann M, Zhuang D, et al. Exenatide once weekly versus twice daily for the treatment of type 2 diabetes: a randomised, open-label, non-inferiority study. *Lancet* 2008; 372: 1240-50.
  15. Victoza (liraglutide injection) Prescribing Information. Available at <http://www.novo-pi.com/victoza.pdf> (accessed 10 December 2014).
  16. Manucci E, Tesi F, Bardini G, Ognibene A, Petracca MG, Ciani S, et al. Effects of metformin on glucagon-like peptide-1 levels in obese patients with or without type 2 diabetes. *Diab Nutr Metab* 2004; 17: 336-342.
  17. Thong KY, Walton C, Ryder REJ. Safety and efficacy of liraglutide 1.2mg in patients with mild and moderate renal impairment: the ABCD nationwide liraglutide audit. *Practical Diabetes* 2013; 30: 71-6.
  18. Jacobsen LV, Hindsberger C, Robson R, Zradkovic M. Effect of renal impairment on the pharmacokinetics of the GLP-1 analogue liraglutide. *Br J Clin Pharmacol* 2009; 68: 898-905.

**Table 1** - Baseline characteristics (continuous variables) and their association with the reporting of gastrointestinal side effects (GISE) and drug discontinuation due to GISE among patients treated with liraglutide

	<i>n</i>	GISE status			<i>P</i> value
		No	Yes, did not discontinue	Yes, discontinued liraglutide	
Age (years)	4213	55 ± 11	55 ± 11	59 ± 12	<0.001
Diabetes duration (years)	3410	9 [6-13]	9 [6-14]	10 [6-14]	0.17
HbA1c (%; mmol/mol)	4153	9.4 ± 1.7; 79 ± 19	9.3 ± 1.6; 78 ± 18	9.4 ± 1.7; 79 ± 19	0.85
Weight (kg)	4262	110.7 ± 22.4	110.0 ± 24.0	106.7 ± 21.0	0.029
BMI (kg/m <sup>2</sup> )	4173	38.9 ± 7.2	38.8 ± 7.4	38.2 ± 7.3	0.41
eGFR* (ml/min/1.73m <sup>2</sup> )	3474	83 ± 21	84 ± 21	77 ± 23	<0.001
ALT (U/L)	2727	28 [21-42]	30 [22-43]	29 [20-44]	0.47
TC (mmol/L)	3486	4.3 ± 1.2	4.2 ± 1.1	4.3 ± 1.0	0.68
HDL-C (mmol/L)	3073	1.1 ± 0.3	1.1 ± 0.3	1.1 ± 0.3	0.80
TG (mmol/L)	2684	2.1 [1.5-2.9]	2.0[1.4,2.9]	2.2 [1.5,3.0]	0.58
SBP (mmHg)	3483	138 ± 18	139 ± 19	141 ± 19	0.15
DBP (mmHg)	3483	79 ± 11	79 ± 11	78 ± 11	0.37

Variables are shown as mean ± sd or median [inter-quartile range]

BMI; body mass index, eGFR; estimated glomerular filtration rate, ALT; alanine transaminotransferase, TC; total cholesterol, HDL-C; high density lipoprotein cholesterol, TG; triglyceride, SBP; systolic blood pressure, DBP; diastolic blood pressure

\*eGFR shown is calculated by the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation. Analyses using the Cockcroft-Gault equation adjusted by body surface area and Modified Diet for Renal Disease (MDRD) equation showed associations with significance of  $P = 0.001$  and  $P = 0.009$ , respectively.

**Table 2** - Baseline characteristics (categorical variables) and their association with the reporting of gastrointestinal side effects (GISE) and drug discontinuation due to GISE among patients treated with liraglutide

	<i>n</i>	GISE status			<i>P</i> value
		No	Yes, did not discontinue	Yes, discontinued liraglutide	
Gender					
Male	4434	2076 (88.9)	144 (6.2)	116 (5.0)	0.13
Female		1823 (86.9)	152 (7.2)	123 (5.9)	
Ethnicity					
Caucasian	3673	2926 (88.1)	224 (6.8)	170 (5.1)	0.88
South-Asian		211 (87.9)	15 (6.3)	14 (5.8)	
Afro-Caribbean		100 (88.5)	9 (8.0)	4 (3.5)	
CKD group*					
1	3452	1297 (89.5)	92 (6.3)	61 (4.2)	0.004
2		1296 (87.8)	88 (6.0)	92 (6.2)	
3		444 (84.4)	37 (7.0)	45 (8.6)	
Metformin use					
No	4442	739 (86.5)	49 (5.7)	66 (7.7)	0.003
Yes		3166 (88.2)	248 (6.9)	174 (4.9)	
Sulphonylurea use					
No	4442	2199 (87.7)	160 (6.4)	148 (5.9)	0.17
Yes		1706 (88.2)	137 (7.1)	92 (4.8)	
TZD use					
No	4442	3141 (87.8)	239 (6.7)	197 (5.5)	0.82
Yes		764 (88.3)	58 (6.7)	43 (5.0)	
Insulin use					
No	4442	2313 (88.6)	179 (6.9)	119 (4.6)	0.012
Yes		1592 (87.0)	118 (6.4)	121 (6.6)	

Variables are shown as n (%)

\*CKD groups shown use eGFR calculated with the CKDEPI equation: CKD Group 1 eGFR >90 ml/min/1.73m<sup>2</sup>, CKD Group 2 eGFR 60-89 ml/min/1.73m<sup>2</sup>, CKD Group 3 eGFR 30-59 ml/min/1.73m<sup>2</sup>. Twenty two patients with eGFR < 30 ml/min/1.73m<sup>2</sup> were excluded due to small numbers in each GISE outcome category. Analyses using the Cockcroft Gault equation adjusted by body surface area and Modified Diet for Renal Disease (MDRD) equation showed associations with significance of *P* < 0.001 and *P* = 0.031, respectively.

TZD; thiazolidinediones

**Table 3** - Ordinal logistic regression analyses of variables associated with gastrointestinal side effects (GISE) or drug discontinuation due to GISE among patients treated with liraglutide

	Model 1		Final model	
	Adjusted OR [95%CI]	<i>P</i> value	Adjusted OR [95%CI]	<i>P</i> value
Insulin use	1.26 [1.02,1.55]	0.035	-	-
Metformin use	0.76 [0.59,0.99]	0.043	0.76 [0.60,0.96]	0.020
Age	1.11 [0.99,1.25]	0.083	1.15 [1.05,1.26]	0.002
Weight	0.96 [0.86,1.06]	0.38	-	-
eGFR	0.98 [0.87,1.10]	0.68	-	-

ORs represent a rise of 1-SD for continuous variable (age, weight, eGFR) and one category change of categorical variable (metformin use, insulin use). Results are sorted by *P* value. Model 1: all significant variables in univariate analyses; Final model: model after sequentially removing variables with the highest non-significant *P* value.

**Figure 1** - Rates of gastrointestinal side effects (GISE) among patients treated with liraglutide, results shown by decades of age. The interaction between age (decades) and GISE outcome was statistically significant ( $P < 0.001$ ). Age 20-29 ( $n=64$ ), age 30-39 ( $n=236$ ), age 40-49 ( $n=862$ ), age 50-59 ( $n=1409$ ), aged 60-69 ( $n=1261$ ), age 70-79 ( $n=339$ ), age 80-89 ( $n=35$ ). Seven patients under 20 years of age were excluded for clarity.

Number of patients	Age (years)	GISE - continued treatment	GISE - discontinued treatment due to GISE	GISE - total
64	20-29	10.9	4.7	15.6
236	30-39	5.5	4.2	9.7
862	40-49	7.4	3.5	10.9
1409	50-59	6.7	4.5	11.2
1261	60-69	7.1	6.0	13.1
339	70-79	6.8	10.6	17.4
35	80-89	0	20.0	20

